Comprehensive Journal of Science
Volume (10), Issue (39), (March, 2026)
ISSN: 3014-6266

مجلة العلوم الشاملة
المجلد (10) العدد 39 مارس 2026
ردمد: 3014-6266

Theoretical Foundations and Properties of Riemann-Liouville Fractional Integration

Intisar Abdullah Aneer

Faculty of Engineering /university of Sabratha

Email:intisar.aneer@sabu.edu.ly

تاريخ الاستلام:2026/01/03 –تاريخ المراجعة: 2026/01/21– تاريخ القبول: 2026/01/30– تاريخ للنشر :2026/02/23

Abstract

This paper provides an extensive analytical investigation into the Riemann-Liouville (R-L) fractional integral, a core operator in fractional calculus. By generalizing the classical n-fold integration process to an arbitrary real order $\alpha > 0$, the R-L operator enables the modeling of systems with long-range memory and non-local interactions. This research provides a rigorous derivation from Cauchy's repeated integration formula and explores fundamental, non-classical properties, including linearity, the semigroup property, and its action on diverse function classes. We present detailed mathematical examples demonstrating the emergence of special functions like the Mittag-Leffler function. Furthermore, a comparative analysis between Riemann-Liouville and Caputo formulations is conducted, followed by an exploration of Laplace transforms in the fractional domain. This comprehensive study aims to bridge the gap between abstract mathematical theory and applied computational modeling, providing a robust framework for understanding fractional-order dynamics .

Keywords: Fractional Calculus, Riemann-Liouville Integral, Caputo Derivative, Mittag-Leffler Function, Non-local Operators, Mathematical Modeling.

Introduction 1

Fractional calculus has emerged as a transformative field of mathematical study ,

generalizing the traditional notions of integer-order differentiation and integration to arbitrary orders. While the history of this field dates back to the late 17th

century with the correspondence between Leibniz and L'Hôpital, it is only in recent

decades that its full potential has been realized in modern science and engineering

[2, 1]. Among the various formulations within this field, the Riemann-Liouville

(R-L) fractional integral stands as the most fundamental operator .

The intrinsic value of the R-L operator lies in its ability to describe phenomena with ”hereditary properties.” Unlike classical calculus, where the derivative or

integral at a point depends only on the immediate neighborhood, fractional operators are non-local. This means they account for the history of the system, making

them indispensable in fields such as anomalous diffusion, signal processing, and

the modeling of complex biological membranes [3,7] .

Despite its theoretical robustness, significant barriers remain in its practical

application. A critical review of existing literature reveals a persistent gap: many

foundational texts focus primarily on direct applications while overlooking detailed explorations of the operator's unique, non-classical mathematical properties. This paper addresses this gap by evaluating the theoretical importance of

the R-L operator and contrasting it with the Caputo fractional integral to delineate

.clear criteria for their respective application .

The remainder of this paper is organized as follows: Section 2 provides the formal mathematical definitions; Section 3 analyzes the linearity and transformation

properties; Section 4 presents a comparative study with the Caputo integral; and

Section 5 concludes with applications and future outlook .

Mathematical Definitions and Notations 2

The Gamma Function 2.1

The foundation of fractional calculus is the Gamma function, Γ(z), which serves as the bridge from discrete factorials to continuous analysis. For any Re(z) > 0, it is defined as [4]:

$$\Gamma(z) = \int_0^\infty e^{-x} x^{z-1} dx \quad (1)$$

The property Γ(z + 1) = zΓ(z) is essential for defining the coefficients of the fractional integral kernel. Without this function, the generalization of (n − 1)! in the Cauchy formula would be impossible .

The Riemann-Liouville Operator 2.2

The Riemann-Liouville fractional integral of order α > 0 for a function f(t) is

$$J_a^\alpha = \frac{1}{\Gamma(\alpha)} \int_a^t (t-\tau)^{\alpha-1} f(\tau) d\tau , \quad t > a$$

This definition can be viewed as an integral transform where the kernel is a power function. The parameter a represents the starting point of the ”memory” of the process .

Analytic Properties and Rigorous Proofs 3

3.1The Linearity Property (Theorem 3.1)

Theorem 3.1. The R-L fractional integral J_a^α is a linear operator on the space $L^1[a, b]$.

Proof. Consider two functions f, g and constants c_1 , c_2. From the definition :

$$J_a^\alpha [c_1 f(t) + c_2 f(t)] = \frac{1}{\Gamma(\alpha)} \int_a^t (t-\tau)^{\alpha-1} [c_1 f(\tau) + c_2 g(\tau)] d\tau$$

$$= \frac{c_1}{\Gamma(\alpha)} \int_a^t (t-\tau)^{\alpha-1} f(\tau) d\tau + \frac{c_2}{\Gamma(\alpha)} \int_a^t (t-\tau)^{\alpha-1} g(\tau) d\tau$$

$$= c_1 j_a^\alpha f(t) + c_2 j_a^\alpha g(t)$$

This confirms that the fractional integral obeys the principle of superposition, a critical requirement for linear system analysis [2] .

3.2The Semigroup Property (Theorem 3.2)

Theorem 3.2. For α, β > 0, the composition of two fractional integrals satisfies

$$J_a^\alpha J_a^\beta f(t) = J_a^{\alpha+\beta} f(t)$$

Proof. Applying the definition twice and utilizing Dirichlet's formula for changing the order of integration [1]:

$$J_a^\alpha \left(J_a^\beta f(t) \right) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^t (t-\tau)^{\alpha-1} \left(\int_a^\tau (\tau - s)^{\beta-1} f(s) ds \right) d\tau$$

By reversing the order of integration and using the substitution $u = (\tau - s)/(t - s)$ the inner integral becomes a Beta function $B(\alpha, \beta)$, which simplifies to the Gamma-based coefficient of $J_a^{\alpha+\beta}$.

3.3Laplace Transform of Fractional Integrals

The Laplace transform is the most common tool for solving fractional differential equations. Let $\mathcal{L}\{f(t)\} = F(s)$. The transform of the R-L integral (for a = 0) is given by :

$$(3) \mathcal{L}\{J_0^\alpha f(t)\} = s^{-\alpha} F(s)$$

This result generalizes the classical rule $\mathcal{L}\{\int f\} = s^{-1} F(s)$.

Comparative Analysis: Riemann-Liouville vs. Caputo 4

4.1 Structural and Conceptual Differences

The comparison between Riemann-Liouville (R-L) and Caputo operators is fundamental to choosing the right mathematical model for a physical system [6] .

1. Mathematical Sequence: D_{RL}^{α}integrates then differentiates, whileD_{Caputo}^{α}

differentiates then integrates. This non-commutativity is a key non-classical

property .

Initial Conditions: Solving $FDEs$ with R-L requires initial values of the form**2.**

which is lack physical meaning . Caputo allows using $f(0), f'(0)$, $[J^{\alpha-1}f\alpha](0) = c$, which represent position and velocity .

5 Applications and Final Conclusions

5.1 Fractional Differential Equations (FDEs)

Example 5.1 (Fractional Relaxation) . Consider $D^{\alpha}y(t) + \lambda y(t) = 0$. The $R - L$ solution involves :

$$y(t) = t^{\alpha-1}E_{\alpha,\alpha}(-\lambda t^{\alpha})$$

where$E_{\alpha,\beta}(z)$denotes the generalized Mittag-Leffler function defined by the following power series:

$$E_{\alpha,\beta}(z) = \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(\alpha k + \beta)}$$

This captures the ”slow” decay observed in complex polymers[7].

6 Conclusion

The Riemann-Liouville operator remains a cornerstone of fractional calculus

This research has successfully clarified its non-classical properties and established

a clear distinction from the Caputo formulation. By providing rigorous proofs for

linearity and the semigroup property, this study provides the necessary groundwork for future developments in non-local dynamics and complex system modeling [9و8] .

References

[1] **Baleanu, D., Diethelm, K., Scalas, E., & Trujillo, J. J. (2012).** *Fractional Calculus: Models and Methods for Non-local Fractional Dynamics*. Singapore: World Scientific Publishing.

[2] **Diethelm, K. (2010).** *The Analysis of Fractional Differential Equations: An Application-Oriented Exposition Using Differential Operators of Caputo Type*. Berlin: Springer-Verlag.

[3] **Hilfer, R. (2000).** *Applications of Fractional Calculus in Physics*. Singapore: World Scientific Publishing.

[4] **Karniadakis, G. E., et al. (2021).** "Physics-informed machine learning." *Nature Reviews Physics*.

[5] **Kilbas, A. A., Srivastava, H. M., & Trujillo, J. J. (2006).** *Theory and Applications of Fractional Differential Equations*. Amsterdam: Elsevier Science B.V..

[6] **Li, C., & Zeng, F. (2020).** *Numerical Methods for Fractional Calculus*. CRC Press.

[7] **Mainardi, F. (2010).** *Fractional Calculus and Waves in Linear Viscoelasticity: An Introduction to Mathematical Models*. London: Imperial College Press.

[8] **Miller, K. S., & Ross, B. (1993).** *An Introduction to the Fractional Calculus and Fractional Differential Equations*. New York: John Wiley & Sons.

[9] **Oldham, K. B., & Spanier, J. (1974).** *The Fractional Calculus: Theory and Applications of Differentiation and Integration to Arbitrary Order*. San Diego: Academic Press.

[10] **Podlubny, I. (1998).** *Fractional Differential Equations: An Introduction to Fractional Derivatives, Fractional Differential Equations, to Methods of Their Solution and Some of Their Applications*. San Diego: Academic Press.

[11] **Samko, S. G., Kilbas, A. A., & Marichev, O. I. (1993).** *Fractional Integrals and Derivatives: Theory and Applications*. Yverdon: Gordon and Breach Science Publishers.

[12] **Sun, H. G., et al. (2022).** "A review of fractional calculus in continuum mechanics." *Journal of Mechanics*.

[13] **Tarasov, V. E. (2023).** *Fractional Dynamics: Applications in Physics and Biology*. De Gruyter.

.