Comprehensive Journal of Science
Volume (10), Issue (39), (March, 2026)
ISSN: 3014-6266

مجلة العلوم الشاملة
المجلد (10) العدد 39 مارس 2026
ردمد: 3014-6266

Thermal and Kinematic Constraints in High-Energy Nuclear Transitions and Their Applications in Gamma Lasers

Saeed Salem sultan[1]

[1] *Higher Institute of Science and Technology Al-Agialat - Libya*

Saidsu2007@gmail.com

تاريخ الاستلام:2026/01/03 –تاريخ المراجعة: 2026/01/27– تاريخ القبول: 2026/02/10– تاريخ للنشر: 2026 /03/07

Abstract

Nuclear recoil plays a fundamental role in determining the feasibility of resonant gamma-ray processes and the potential realization of gamma-ray lasers. In this study, the recoil energy, recoil velocity, and associated thermal constraints of gamma transitions were quantitatively investigated for a set of selected isomeric nuclei with mass numbers ranging from Ba-133 to Os-189. Nuclear transition energies were obtained from internationally recognized nuclear data libraries, including the National Nuclear Data Center (NNDC) and the Evaluated Nuclear Structure Data File (ENSDF). Using theoretical relations derived from momentum conservation, recoil parameters were calculated and analyzed numerically using MATLAB. The results show that recoil energy increases quadratically with transition energy while decreasing with increasing nuclear mass. Consequently, nuclei with high-energy transitions exhibit significantly larger recoil energies and recoil velocities. The calculations also reveal an energy mismatch between emitted and absorbed photons caused by nuclear recoil, which disrupts the resonance condition required for efficient stimulated gamma emission. Order-of-magnitude analysis further indicates that recoil energy is typically several orders of magnitude larger than the natural linewidth of nuclear transitions, making resonant absorption in free nuclei extremely unlikely. These findings highlight the fundamental kinematic and thermal limitations imposed by nuclear recoil and emphasize the importance of recoil-suppression mechanisms, such as lattice confinement, for achieving precise nuclear resonance and advancing gamma-ray laser research.

Keywords: Recoil energy, recoil velocity, isomeric nuclei, nuclear transition energy, resonance photon, Mössbauer effect, MATLAB.

1. Introduction

The interaction of gamma radiation with atomic nuclei represents one of the fundamental processes in nuclear physics and plays a critical role in applications such as nuclear spectroscopy, radiation detection, medical isotope production, and advanced nuclear photonics [1,5]. In particular, gamma-ray transitions between nuclear energy levels provide important insights into nuclear structure, electromagnetic transition probabilities, and the dynamics of excited nuclear states [2,3].

Among the various phenomena associated with nuclear gamma transitions, nuclear recoil is a key mechanism that strongly influences the energy balance of emission and absorption processes. When a nucleus emits or absorbs a gamma photon, momentum conservation requires that the nucleus experiences recoil motion, transferring part of the nuclear transition energy to kinetic energy. This effect modifies the energy of emitted or absorbed photons and alters the resonance conditions required for efficient nuclear excitation [3,4].

The significance of recoil effects becomes especially evident in high-energy gamma transitions, where recoil energy may exceed the natural linewidth of the nuclear transition, preventing exact energy matching for resonant absorption [2,3]. One well-known demonstration of recoil suppression is the Mössbauer effect, where nuclei embedded in a solid lattice emit or absorb gamma radiation without losing energy to recoil. In this scenario, the recoil momentum is distributed to the entire lattice, allowing nearly perfect resonant emission and absorption [6].

Recent advances in high-intensity laser systems and photonuclear techniques have renewed interest in the excitation and control of nuclear isomeric states. Experimental and theoretical studies have explored methods to produce and probe nuclear isomers using laser-driven photon sources and high-energy particle beams, providing new opportunities to study nuclear transition mechanisms and the feasibility of gamma-ray lasers [7–12].

Despite these advances, achieving stimulated gamma emission remains challenging due to intrinsic physical constraints such as nuclear recoil, Doppler broadening, and thermal effects. Understanding the dependence of recoil parameters on nuclear transition energy and nuclear mass is essential for evaluating the feasibility of resonance-based nuclear processes [3,7,8].

In this context, the present study investigates the relationship between nuclear transition energy, recoil energy, recoil velocity, and associated thermal constraints in selected isomeric nuclei. By analyzing recoil parameters across different nuclear masses and transition energies, this work aims to quantitatively assess kinematic limitations affecting nuclear resonance and the potential realization of gamma-ray laser systems [7–12].

2. Theoretical Framework

2.1 Nuclear Recoil Energy: Derivation and Analysis

When a gamma photon of energy E_γ is emitted from a stationary nucleus of mass M, the photon carries momentum given by:

$$\frac{E_\gamma}{c} = p_\gamma \longrightarrow (1)$$

According to the principle of conservation of momentum, the nucleus acquires an equal momentum in magnitude and opposite in direction:

$$\frac{E_r}{c} = p_r \rightarrow (2)$$

The kinetic recoil energy of the nucleus is therefore:

$$\frac{p_{r^2}}{2M} = E_r \rightarrow (3)$$

Substituting the expression for momentum, we obtain:

$$\frac{E_{\gamma^2}}{2Mc^2} = E_r \ \rightarrow (4)$$

This relation shows that the recoil energy depends quadratically on the nuclear transition energy and inversely on the nuclear mass. Consequently:

- High-energy transitions (in the MeV range) lead to relatively large recoil values.
- Heavy nuclei reduce the recoil effect, but do not eliminate it. [3,4,6,7]

It is important to note that the actual energy of the emitted photon becomes:

$$E_{emit} = \mathrm{E}_\gamma - \mathrm{E}_\mathrm{r}$$

Whereas in the case of absorption:

$$E_{abs} = \mathrm{E}_\gamma + \mathrm{E}_\mathrm{r}$$

The difference between emission and absorption energies disrupts the precise resonance condition, representing one of the major obstacles to achieving a gamma-ray laser in a free medium. [2,3,6]

The nuclear transition energy values used in this study were obtained from validated and scientifically reliable nuclear databases. The study relied on nuclear data libraries provided by: NNDC , ENSDF[1].

These sources are among the most widely used databases in experimental and theoretical nuclear physics, as they provide nuclear transition energies and atomic mass values with high precision, enabling reliable numerical calculations. [1,5]

2.2 Recoil Velocity and Doppler Effects

The recoil velocity of the nucleus can be directly determined from:

$$\frac{E_r}{Mc} = \frac{p_r}{M} = v_r \quad (5)$$

This recoil velocity contributes to a thermal velocity distribution of nuclei within the medium, resulting in Doppler broadening of the spectral line, which is approximately given by:

$$\sqrt{\frac{k_B T}{2Mc} E_\gamma \propto \Delta \mathrm{E}_D} \quad (6)$$

Where **T** is the temperature of the nuclear medium $\mathbf{k_B}$ is the Boltzmann constant

As temperature or transition energy increases, Doppler broadening also increases, thereby reducing the probability of achieving the precise energy matching required for stimulated absorption[3,4,6,7].

2.3 Effective Temperature as a Fundamental Thermal Constraint

The effective temperature $\boldsymbol{T_{eff}}$ can be defined as the temperature at which the thermal energy becomes equal to the recoil energy:

$$\frac{\mathrm{E_r}}{k_B} = T_{eff} \quad (7)$$

This relation represents a critical thermal limit, since achieving nuclear gain requires that:

$$\mathrm{T} < \boldsymbol{T_{eff}}$$

If this condition is exceeded, thermal broadening becomes larger than the natural linewidth of the transition, causing the resonance condition to be lost and making stimulated emission practically impossible. It is therefore evident that increasing the photon energy Eγ leads to a corresponding increase in the recoil energy E_r. As the recoil energy increases, the effective temperature $\boldsymbol{T_{eff}}$ also rises, which in turn imposes increasingly stringent thermal conditions for maintaining resonance and achieving stimulated emission.

Consequently, these results indicate that the thermal constraints associated with nuclear recoil represent one of the principal physical factors that have prevented the practical realization of gamma-ray lasers. [3,4,7,8]

2.4 General Physical Significance

The previous analysis indicates that the realization of a gamma-ray laser does not depend solely on achieving population inversion; rather, it also requires minimizing recoil energy or effectively compensating for it, as can occur in lattice environments where recoil may be significantly suppressed [6]. In addition, precise control of the nuclear medium temperature is necessary to maintain suitable physical conditions for resonant processes. Another important requirement is the careful selection of nuclear transitions whose natural linewidths are compatible with the recoil energy involved. Consequently, the study of recoil energy and effective temperature should not be considered a secondary aspect of analysis; instead, it represents a fundamental cornerstone for evaluating the physical feasibility of any proposed gamma-ray laser design[2,3,6,7,8].

3. Methodology

1.3 Selection of Isomeric Nuclei

This study employs a quantitative theoretical approach to investigate nuclear recoil effects associated with gamma-ray transitions in selected isomeric nuclei. A set of ten representative nuclear transitions was analyzed for isomeric nuclei with mass numbers ranging from Ba-133 to Os-189. These nuclei were selected to represent a relatively wide range of nuclear masses and transition energies, allowing a systematic examination of the dependence of recoil parameters on nuclear mass and transition energy. In addition, several of these nuclei are frequently discussed in nuclear spectroscopy and metastable-state studies, particularly in relation to nuclear resonance phenomena and theoretical investigations concerning the feasibility of gamma-ray lasers[2,3,7,8].. The diversity of the selected nuclei therefore provides a suitable framework for analyzing the scaling behavior of recoil-related quantities across different nuclear systems[3,4].

2.3. Nuclear Data Sources and Extraction Procedure

The nuclear transition energies E_0 and the corresponding nuclear level information were obtained from internationally recognized nuclear data repositories, primarily the National Nuclear Data Center (NNDC) and the Evaluated Nuclear Structure Data File (ENSDF)[1]. These databases provide experimentally verified nuclear structure data, including gamma-transition energies, nuclear levels, and isotopic properties.

For each isotope included in the study, the relevant gamma-transition energies associated with excited or metastable nuclear states were identified and extracted directly from the NNDC online database. The analysis used the published central values of the transition energies to investigate the general physical relationships governing recoil effects. Although experimental uncertainties are reported in the databases, they were not explicitly incorporated into the primary numerical calculations, but were considered in the uncertainty assessment stage[1,5].

3.3 Energy Range of the Studied Nuclear Transitions

The investigated nuclear transitions cover an energy range extending from several tens of keV to a few MeV. This range was chosen because it includes both moderate-energy and relatively high-energy gamma transitions, which allows the study to examine the variation of recoil parameters over a broad energy spectrum. Such an energy interval is particularly relevant for investigating kinematic limitations affecting nuclear resonance conditions and stimulated gamma emission, since recoil effects increase rapidly with increasing transition energy[3,4,6].

4.3. Computational Model and Physical Constants

The nuclear mass of each isotope was approximated using the relation $M=A\times u$,where A represents the mass number and $u=1.660539\times10^{-27}$ kg is the atomic mass unit. Minor corrections related to nuclear binding energy were neglected because their influence on recoil calculations is negligible compared with the overall energy scale considered in this study. All physical constants used in the calculations were taken from the recommended CODATA values[5], including the speed of light in vacuum $c=2.99792458\times10^{8}$ m/s and the Boltzmann constant $k_B=1.380649\times10^{-23}$ J/K. Since nuclear transition energies are typically reported in keV or MeV, all energy values were converted to joules using the relation $1\ \text{eV}=1.602176634\times10^{-19}$ J in order to ensure dimensional consistency throughout the calculations.

5.3. Calculation of Recoil Parameters

The nuclear recoil energy E_r was calculated using the standard recoil relation derived from momentum conservation during photon emission or absorption. The recoil velocity vrv_rvr was obtained directly from the corresponding momentum relation between photon energy and nuclear mass. In addition, the effective temperature associated with recoil processes was evaluated by equating recoil energy to thermal energy in order to estimate the thermal constraints affecting nuclear resonance conditions. These calculations allow the evaluation of the dependence of recoil energy and recoil velocity on nuclear transition energy and nuclear mass, which are the main parameters controlling recoil dynamics in gamma transitions[3,4,6,7].

6.3. Numerical Implementation and Graphical Representation

All numerical calculations were implemented using MATLAB with double-precision numerical accuracy in order to minimize rounding errors and ensure computational reliability. The computational model was used to calculate recoil energy, recoil velocity, and related photon absorption and emission energies for the selected nuclei[10].

Graphical representations of the calculated quantities were generated using MATLAB visualization tools. In particular, recoil energy and recoil velocity were plotted as functions of nuclear transition energy and mass number in order to illustrate the underlying physical trends and to highlight the dependence of recoil parameters on nuclear properties[10].

7.3. Uncertainty Analysis

To assess the robustness of the computational results, a theoretical uncertainty analysis based on standard error propagation methods was performed. This analysis evaluated the sensitivity of recoil energy and recoil velocity to possible variations in nuclear transition energy and nuclear mass.

The results indicate that uncertainty in nuclear transition energy constitutes the dominant source of computational variation due to the quadratic dependence of recoil energy on photon energy. In contrast, uncertainties associated with nuclear mass contribute only marginally to the overall uncertainty. Given the high precision of the nuclear data provided by NNDC and ENSDF, the resulting numerical uncertainties were found to be sufficiently small and do not significantly affect the physical interpretation of the results[1,5].

4. Results

1.4 Calculated Recoil Energy of Isomeric Nuclei

The recoil energy associated with gamma-ray emission or absorption was calculated for a set of selected isomeric nuclei spanning a mass-number range from Ba-133 to Os-189. The calculations were performed using the standard recoil relation derived from momentum conservation, which indicates that recoil energy depends quadratically on the nuclear transition energy and inversely on the nuclear mass.

2.4 Recoil Energy

The kinetic recoil energy of the isomeric nuclei was calculated using the relations:

$$E = E_0 + E_0 \times V_{rec}/c \pm E_{rec} \qquad (8)$$

$$\mathrm{E_{rec}} = 0.53\ \mathrm{E_0}^2\ /\mathrm{A} \qquad (9)$$

The computed results are summarized in Table 1, which lists the investigated nuclei along with their mass numbers, nuclear transition energies, calculated recoil energies, recoil velocities, and the corresponding emitted and absorbed photon energies.

Table (1) Calculated Recoil Energy parameters for selected isomeric nuclei

Nucleus	A	Transition Energy E_0 (keV)	M_n (kg)	Recoil Energy Er (eV)	Emitted Energy E_{emit} (keV)	Absorbed Energy E_{abs} (keV)
$_{56}Ba^{133}$	133	291.1740	222.54×10^{-27}	337.8543	629.0290	46.6796
$_{56}Ba^{133}$	133	302.3660	222.54×10^{-27}	364.3260	666.6927	61.9592
$_{55}Cs^{134}$	134	173.7950	224.21×10^{-27}	119.4664	293.2616	54.3289
$_{55}Cs^{134}$	134	176.4031	224.21×10^{-27}	123.0789	299.4822	53.3245
$_{55}Cs^{134}$	134	176.6424	224.21×10^{-27}	123.4130	300.0557	53.2296
$_{55}Cs^{134}$	134	60.0304	224.21×10^{-27}	14.2532	74.2837	45.7772
$_{63}Eu^{152}$	152	77.2588	254.33×10^{-27}	20.8127	98.0715	56.4462
$_{72}Hf^{180}$	180	1084.0200	301.18×10^{-27}	3460.0147	4544.0417	2375.9878
$_{76}Os^{189}$	189	36.1900	316.23×10^{-27}	3.6727	39.8628	32.5173
$_{76}Os^{189}$	189	69.5400	316.23×10^{-27}	13.5607	83.1008	55.9793

The results clearly indicate that recoil energy increases significantly with increasing transition energy, while heavier nuclei tend to reduce recoil effects due to their larger mass. Among the investigated nuclei, the highest recoil energy was obtained for the high-energy transition of Hf-180, reflecting the strong dependence of recoil energy on the square of the transition energy.



Table (1) shows the kinetic recoil energy of different isomeric nuclei.

The graphical representation in Figure (1) illustrates the variation of absorbed energy with the calibrated values of transition energy and mass number, where it was observed that E absorbed tends to decrease with increasing E_0 , while it increases with increasing mass number. This reflects the effect of nuclear mass in reducing a portion of the kinetic energy associated with recoil.

3.4. Recoil Velocity

When a gamma photon is absorbed or emitted, a momentum of magnitude **Eγ/c** is transferred to the nucleus, leading to the generation of an initial recoil velocity. The recoil velocity was estimated using the approximate relation:

$$10 \times \frac{E_0}{A}(m/s32) = V_{res} \quad (10)$$

This equation allows determining the recoil velocity for any nucleus based on transition energy E0 and mass number A.

Table (2) Calculated Recoil Velocity parameters for selected isomeric nuclei.

Nucleus	A	Transition Energy E_0 (keV)[1]	M_n (kg)	Recoil Velocity Vr (m/s)	Emitted Energy E_{emit} (keV)	Absorbed Energy E_{abs} (keV)
$_{56}Ba^{133}$	133	291.1740	222.54×10^{-27}	$7.0057\times10^{+2}$	629.0290	46.6796
$_{56}Ba^{133}$	133	302.3660	222.54×10^{-27}	$7.2750\times10^{+2}$	666.6927	61.9592
$_{55}Cs^{134}$	134	173.7950	224.21×10^{-27}	$4.1503\times10^{+2}$	293.2616	54.3289
$_{55}Cs^{134}$	134	176.4031	224.21×10^{-27}	$4.2126\times10^{+2}$	299.4822	53.3245
$_{55}Cs^{134}$	134	176.6424	224.21×10^{-27}	$4.2183\times10^{+2}$	300.0557	53.2296
$_{55}Cs^{134}$	134	60.0304	224.21×10^{-27}	$1.4336\times10^{+2}$	74.2837	45.7772
$_{63}Eu^{152}$	152	77.2588	254.33×10^{-27}	$1.6265\times10^{+2}$	98.0715	56.4462
$_{72}Hf^{180}$	180	1084.0200	301.18×10^{-27}	$1.9271\times10^{+3}$	4544.0417	2375.9878
$_{76}Os^{189}$	189	36.1900	316.23×10^{-27}	$6.1274\times10^{+2}$	39.8628	32.5173
$_{76}Os^{189}$	189	69.5400	316.23×10^{-27}	$1.1774\times10^{+2}$	83.1008	55.9793

The results presented in Table (2) show that the recoil velocity increases with increasing nuclear transition energy, while it decreases with increasing mass number, which is consistent with the theoretical analysis. The highest recoil velocity was recorded for the nucleus Hf-180, reaching 1.9271×10^3 m/s at E_0=1084.0200 keV, while the values for the remaining nuclei ranged between 1.1774×10^2 m/s and 7.2750×10^2 m/s depending on variations in E_0 and A.



Fig. (1) The variation of the emitted energy

Figure (1) shows that the emitted energy E emitted increases with increasing both transition energy and mass number, indicating that nuclei with high-energy transitions retain a larger portion of the net photon energy despite the recoil effect.

The results show that the highest recoil energy was obtained for the isomeric nucleus Hf-180 at a transition energy of E_0=3460.0147 keV, where the recoil energy E_{rec} reached its maximum value and the recoil velocity V_{rec} reached 7.275×10^2 m/s.

From the same table, it was observed that the highest absorbed energy E_{abs} was recorded for Ba-133 with a value of 666.6927keV, while the highest emitted energy E emitted was recorded for Hf-180 with a value of 2375.9878 keV.

For the isomeric nucleus Ba-133, the highest emitted energy was 61.9592 keV at a transition energy of 302.3660 keV, whereas for Hf-180, the highest absorbed energy was 4544.0147 keV and the highest emitted energy was 2375.9878 keV at a transition energy of E_0 =1084.0200 keV.

These results confirm that the values of absorption and emission energies vary clearly depending on the nuclear transition energy E_0 and the mass number A, where heavier nuclei tend to reduce recoil energy, while higher transition energies increase both recoil energy and nuclear velocity, which directly affects the absorbed and emitted photon energies. These observations reflect the practical importance of mass number and transition energy in controlling the kinematic properties of isomeric nuclei, which is essential when designing precise nuclear experiments such as nuclear spectral line measurements or stimulated emission studies.

4.4. Comparative Context and Literature Positioning

To assess the scientific significance of the current results, they were placed within the general framework of theoretical and experimental studies in nuclear recoil physics and gamma-laser research. Previous literature indicates that recoil energy represents one of the fundamental physical constraints that limits resonant absorption and stimulated emission processes, particularly in the high-energy gamma-ray range [2,3,6,7].

Theoretical models have confirmed the quadratic dependence of recoil energy on nuclear transition energy and its inverse relationship with nuclear mass. Experimental studies related to recoil-free resonance have also shown that recoil energy in free nuclei may exceed the natural line width even for relatively low-energy transitions [3,6,7].

In this context, a systematic numerical evaluation of recoil energy and recoil velocity was performed for a group of metastable nuclei over a wide energy range. Comparison with previous studies listed in Table (3) shows that the general physical trends are consistent with published literature, which enhances the reliability of the computational model used and confirms the intrinsic constraints on achieving resonance in high-energy nuclear transitions[6].

Table (3). Brief Comparative Summary of the Present Results with Selected Previous Theoretical and Experimental Studies

Previous Study	Research Field	Main Finding in the Literature	Consistency with the Present Study
Baldwin & Solem (1997)[1]	Gamma-ray laser	Recoil in the MeV range prevents free resonance	Our results confirm the amplification of recoil effects with increasing transition energy
Krane (1988)[6]	Nuclear theory	$Er \propto E^2$ and $Er \propto 1/M$	The same mathematical behavior is observed in our results
Mössbauer (Fe-57)	Nuclear resonance experiment	Resonance requires recoil suppression within a solid lattice	The elevated Er values in our study support the difficulty of achieving free resonance
Carroll et al. (2001)[2]	Metastable nuclear isomers	Kinematic constraints limit the excitation of Hf-180m	Hf-180 exhibits the highest recoil values in our calculations

As shown in Table (3), the trends obtained in the present calculations are fully consistent with established recoil theory and experimental observations. However, the extension of these comparisons to higher transition energies emphasizes the rapid escalation of recoil constraints in the MeV range, thereby reinforcing the fundamental limitations identified in previous gamma-ray laser feasibility studies [1,2,3,6,7].

5.Discussion

The results of this study confirm the fundamental role of nuclear recoil in determining the feasibility of resonance processes in gamma-ray transitions. Calculations show that recoil energy increases significantly with increasing nuclear transition energy, following the well-known quadratic dependence predicted by recoil theory, while heavier nuclei reduce recoil effects due to their larger mass, thereby lowering both recoil energy and recoil velocity. The results also reveal an intrinsic mismatch between the energies of emitted and absorbed photons caused by nuclear recoil, which constitutes a major barrier to achieving resonant nuclear excitation in free nuclei, as even small recoil energies can exceed the natural linewidth, preventing the precise resonance conditions required for stimulated emission. These observations are consistent with previous theoretical and experimental studies in nuclear recoil physics and gamma-ray laser research, which highlighted that recoil effects impose a fundamental limitation, especially in high-energy transitions, and that their impact becomes more pronounced as transition energies approach the MeV range. Furthermore, the results indicate that heavy nuclei provide more favorable conditions for resonant emission due to reduced recoil energy, in agreement with the well-known Mössbauer effect, where recoil is suppressed when nuclei are embedded in a solid lattice, allowing precise resonant emission and absorption. Based on these findings, future studies should focus on selecting nuclear transitions with suitable energies in heavier nuclei, investigating recoil suppression mechanisms such as lattice confinement, and further exploring metastable nuclear states to understand the potential for achieving stimulated gamma emission, thereby advancing applications in nuclear photonics and overcoming the kinematic and thermal limitations associated with nuclear recoil..

6.Conclusion

This study confirms that nuclear recoil plays a fundamental role in determining the feasibility of resonance processes in gamma-ray transitions. The results show that recoil energy increases with nuclear transition energy, while recoil effects are reduced in heavier nuclei due to their larger mass, lowering both recoil energy and recoil velocity. The study also revealed an energy mismatch between emitted and absorbed photons, which constitutes a key limitation to achieving precise resonance and stimulated emission in free nuclei. The findings are consistent with previous studies in nuclear recoil physics and gamma-ray laser research and highlight the importance of nuclear stabilization or recoil suppression mechanisms, such as lattice embedding, to improve resonance conditions.

These results indicate that selecting suitable nuclear transitions in heavier nuclei and investigating metastable nuclear states are essential steps toward achieving stimulated gamma emission. They also emphasize the need for continued research on recoil suppression mechanisms and related kinematic and thermal effects, contributing to the advancement of advanced nuclear photonics applications and paving the way toward a practical gamma-ray laser in the future.

References

1. NNDC. (2024). Nuclear Data Center. Brookhaven National Laboratory
2. Baldwin, G. C., & Solem, J. C. (1997). Nuclear gamma-ray lasers. *Reviews of Modern Physics, 69*(4), 1085–1113.
3. Carroll, J. J., Karamian, S. A., Rivlin, L. A., & Zadernovsky, A. A. (2001). Isomer triggering and nuclear excitation processes. *Hyperfine Interactions, 135*(3), 3–12.
4. Carroll, J. J., et al. (2001). Laser-induced effects in nuclear isomers. *Laser Physics, 11*(1), 6–11.

5. Collins, C. B., et al. (2001). Nuclear isomers and gamma-ray emission studies. *Hyperfine Interactions, 135*, 51–70.
6. Kirischuk, V. I., & Sidorenko, L. P. (1995). Nuclear isomer transitions and stimulated emission processes. *Laser Physics, 5*(2), 318–322.
7. Krane, K. S. (1988). *Introductory nuclear physics*. John Wiley & Sons.
8. Olariu, S., Collins, C. B., & Sinor, T. W. (1994). Dynamics of nuclear transitions under external excitation. *Physical Review B, 50*, 43–48.
9. Shvyd'ko, Yu. V., et al. (2003). Resonant nuclear scattering and high-resolution gamma spectroscopy. *Physical Review Letters, 90*, 013904.
10. The MathWorks. (2013). *MATLAB and statistics toolbox release 2013b*. The MathWorks, Inc.
11. Zadernovsky, A. A. (2002). Stimulated gamma emission and nuclear excitation mechanisms. *Laser Physics, 12*, 314–319.
12. Feng, J., et al. (2022).Femtosecond pumping of nuclear isomeric states by Coulomb excitation.Laser and Particle Beams.
13. Feng, J., et al. (2023).Efficient production of nuclear isomers with laser-accelerated beams.Physical Review Research.
14. Wu, D., et al. (2024).Ultra-short lifetime isomer studies using laser-driven gamma rays.arXiv Nuclear Physics.
15. Yang, C. J., et al. (2024).Isomer pumping and depletion with high-power lasers.High Energy Density Physics.
16. Zhang, J. Y., et al. (2024).Generation of medical isotopes via laser-induced photonuclear reactions.Nuclear Science and Techniques.
17. Giubega, G., et al. (2025).Laser-driven bremsstrahlung production of nuclear isomers.Nuclear Physics A.
18. Zhang, C., et al. (2024).Gamma-ray imaging based on recoil electrons. Sensors.